I49.89:82/47-1/map E4

MMS 85-0011
FWS/OBS-82/47.1

MAP E4

Fort Walton Beach

FLORIDA ECOLOGICAL ATLAS

HYDROLOGY AND CLIMATOLOGY





TOPICAL MAPS IN THIS ATLAS:

A. BIOLOGICAL RESOURCES
B. SOCIOECONOMIC FEATURES
C. SOILS AND LANDFORMS
D. OIL, GAS AND MINERALS
E. HYDROLOGY AND CLIMATOLOGY

1984

Produced by U.S. Fish and Wildlife Service and Minerals Management Service

Projection and 10,000 meter grid, zone 17: Universal Transverse Mercator
25,000-foot grid ticks based on Florida coordinate system, west zone

Overlay material compiled in 1983 from latest available data obtained from





Due to coastline configuration the inundation zones represent generalized boundaries.

Produced by U.S. Fish and Wildlife Service and Minerals Management Service

Projection and 10,000 meter grid, zone 17, Universal Transverse Mercator
20,000-foot grid ticks based on Florida coordinate system, west zone

Overlay material compiled in 1983 from latest available data obtained from U.S. Geological Survey environment agencies and other sources.

G U L F O F M E X I C O

—50'— POTENTIOMETRIC CONTOURS OF THE FLORIDAN AQUIFER
The map represents dry season water levels present in rights cased wells during the period 15 May through 5 June of 1980. The contour interval is 5 and 10 feet. Contour lines have been generalized to allow for non-simultaneous water level measurement throughout the area.
Groundwater flow is downgradient from potentiometric highs and perpendicular to potentiometric contour lines.
(Measured in feet above mean sea level)

••••52'•••• MEAN ANNUAL ISOHYETAL (RAINFALL) CONTOURS
Based on 1980 National Weather Service Normal Base Period)
The contour interval is 1.0" (refer to narrative for monthly isohyetal maps and for trend evaluation)

6 NATIONAL WEATHER SERVICE 30-YEAR CLIMATOLOGICAL STATION
(Number in the box on map refers to station name in supplemental legend

WIND ROSE STATION
Each wind rose represents 5% frequency interval. The average wind speed (mph) is based on the

NATIONAL WEATHER SERVICE 30-YEAR CLIMATOLOGICAL STATION
Number on map refers to station in explanatory legend

WIND ROSE STATION
Concentric circles represent 5% frequency of occurrence. Shaded speed (mph) in force in the octant quadrant of each vector direction (Wind rose data represents wind data that has been summarized by the National Climatic Center, Asheville, North Carolina, Naval Air Stations and WMO Offices)

OFFSHORE SURFACE CURRENT ROSE
Resultant vectors for each season (Each vector represents the resultant vector in proportion to the velocity (Concentric circles represent 3 centimeters per second)
Vectors defined by the New England Coastal Engineers Computer Model G.N.I.
S Spring
Su Summer
F/W Fall/Winter

HURRICANE INUNDATION ZONES
(Maximum areas subject to flooding by Storm Category, as defined by National Weather Service Computer Models)

Category #1 Storm (74-95 mph winds; probability of occurrence 1-25 years)

Category #2 Storm (96-110 mph winds; probability of occurrence 1-27 years)

Category #3 Storm (111-130 mph winds; probability of occurrence 1-42 years)

Category #4 Storm (131-155 mph winds; probability of occurrence no historical record)

Category #5 Storm (+ 155 mph winds; probability of occurrence no historical record)

(Hurricane inundation zones on this map represent generalized areas of maximum areal extent of flooding from storm surge). The scale and detail of this map are adequate for planning purposes only. This map is unsuitable for determining the vulnerability of individual buildings to storm surge)

U.S. GEOLOGICAL SURVEY

THE FLORIDA ECOLOGICAL ATLAS WAS PRODUCED BY THE U.S. FISH AND WILDLIFE SERVICE DIVISION OF BIOLOGICAL SERVICES, NATIONAL COASTAL ECOSYSTEMS TEAM, AS A COOPERATIVE EFFORT WITH THE FLORIDA DEPARTMENT OF COMMUNITY AFFAIRS, REGIONAL PLANNING COUNCILS, WATER MANAGEMENT DISTRICTS, LOCAL PLANNING AGENCIES, AND THE GENERAL PUBLIC TO DETERMINE AND DOCUMENT THE ECOLOGICAL RESOURCES, HABITAT REQUIREMENTS, AND FISH AND WILDLIFE REQUIREMENTS, AND THE MAINTENANCE OF ECOLOGICAL DIVERSITY. THE STUDY AREA INCLUDES THE COASTAL COUNTIES: ESCAMBIA, SANTA ROSA, OKALOOSA, WALTON, BAY, GULF, FRANKLIN, WAKULLA, JEFFERSON, TAYLOR, DIXIE, LEVY, CITRUS AND HERNANDO, AND EXTENDS INLAND APPROXIMATELY 10 MILES. THERE IS NO ATTEMPT, IN THE DESIGN OF THIS ATLAS, TO ALTER OR AFFECT IN ANY WAY THE STATUTORY JURISDICTION OF ANY FEDERAL, STATE, OR LOCAL GOVERNMENT TO ESTABLISH AND ENFORCE LAWS, RULES, REGULATIONS, OR POLICIES OF THEIR AGENCIES.

THIS MAP WAS BEEN COMPILED FROM A VARIETY OF SOURCE MATERIALS INCLUDING MAPS, AERIAL PHOTOS, AND REPORTS. SINCE MANY SOURCES WERE USED, THERE MAY BE SOME DISCREPANCIES IN THE LOCATION OF MAPPED FEATURES. THE USER SHOULD NOT RELY ON THESE MAPS FOR PRECISE MEASUREMENTS OF DISTANCES OR AREAS.

PRIMARY SOURCES OF MAPPED INFORMATION:

HYDROLOGY: Potentiometric Surface of the Floridan Aquifer, Northwest Florida Water Management District, Information Publication 83-1, 1983.
Potentiometric Surface of the Floridan Aquifer, Suwannee River Water Management District, Open File Report, 1982.
Potentiometric Surface of the Floridan Aquifer, Southwest Florida Water Management District, Brooksville, Florida, 1982.
Rainfall Intensities for Selected Design Data from Hurricanes, SEA-SWH-Note, National Oceanic and Atmospheric Administration, Silver Spring, Maryland, 1978.
Surface Water Supply of the United States, 1961-65, Geological Survey Professional Paper 1818, 1967.
Water Quality Inventory for the State of Florida, 1982 305(b) Technical and Publication, Tallahassee, Florida, 1982.
Current Atlas of the West Florida Shelf, New England Coastal Engineers, Bangor, Maine, 1982.

CLIMATOLOGY: National Climatic Center, Asheville, North Carolina, 1982.

FOR COMPLETE REFERENCES SEE MAP NARRATIVE

SCALE 1:100 000

KILOMETERS 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39